

RECONSTRUCTING COMPOSITIONS

VINCENT VATTER*

School of Mathematics and Statistics

University of St Andrews

St Andrews, Fife, Scotland

vince@mcs.st-and.ac.uk

<http://turnbull.mcs.st-and.ac.uk/~vince>

We consider the problem of reconstructing compositions of an integer from their subcompositions, which was raised by Raykova (albeit disguised as a question about layered permutations). We show that every composition w of $n \geq 3k + 1$ can be reconstructed from its set of k -deletions, i.e., the set of all compositions of $n - k$ contained in w . As there are compositions of $3k$ with the same set of k -deletions, this result is best possible.

Introduction. The Reconstruction Conjecture states that given the multiset of isomorphism types of 1-vertex deletions (briefly, *1-deletions*) of a graph G — the *deck* of G — on three or more vertices, it is possible to determine G up to isomorphism. The stronger set version of the conjecture due to Harary [5] only allows access to the *set* of 1-deletions and requires G to have four or more vertices. These conjectures can be made even more difficult by considering k -deletions instead of 1-deletions, for which we refer to Manvel [7].

Such reconstruction questions extend naturally to other combinatorial contexts. For example, Schützenberger and Simon (see Lothaire [6, Theorem 6.2.16]) proved that every word of length $n \geq 2k + 1$ can be reconstructed from its set of k -deletions (i.e., subwords of length $n - k$). This bound is tight because the words $(ab)^k$ (the word with ab repeated k times) and $(ba)^k$ have the same set of k -deletions: all words of length k over the set $\{a, b\}$. Answering a question of Cameron [4], Pretzel and Siemons [8] considered the partition context, where they proved that every partition of $n \geq 2(k + 3)(k + 1)$ can be reconstructed from its set of k -deletions. (This bound is not known to be tight.)

Motivated by a question of Raykova [9] (described at the end of the paper), we consider the problem of set reconstruction for compositions (ordered partitions), establishing the following result.

*Supported by EPSRC grant GR/S53503/01.

Date: February 2, 2008

Key words and phrases. composition, layered permutation, reconstruction

AMS 2000 Subject Classification. 68R15, 06A07, 05A17, 05C60

Theorem 1. *All compositions of $n \geq 3k + 1$ can be reconstructed from their sets of k -deletions.*

Our proof of Theorem 1 illustrates an algorithm to perform the reconstruction. Perhaps more convincing than the proof is the Maple implementation of this algorithm, available from the author's homepage.

Notation. We view a composition as a word w whose letters are positive integers, i.e., a word in $\mathbb{P}^*$. We denote the length of w by $|w|$ and the sum of the entries of w by $\|w\|$, and say that w is a composition of $\|w\|$. A 1-deletion of w is a composition that can be obtained either by lowering a ≥ 2 entry of w by 1 or by removing an entry of w that is equal to 1. A 2-deletion is then a 1-deletion of a 1-deletion, and so on.

This notion naturally defines a partial order[†] on compositions: $u \leq w$ if w contains a subword $w(i_1)w(i_2) \cdots w(i_\ell)$ of length $\ell = |u|$ such that $u(j) \leq w(i_j)$ for all $1 \leq j \leq \ell$. (We refer to the indices $i_1 < \cdots < i_\ell$ as an *embedding* of u .) For example, $1211 \leq 21312$ because of the subword 2312 . If $u \leq w$ then u is a $(\|w\| - \|u\|)$ -deletion of w . Returning to the previous example, $\|21312\| = 9$ and $\|1211\| = 5$, so 1211 is a 4-deletion of 21312 .

A lower bound. In the context of words, the fact that the sets of k -deletions of $(ab)^k$ and $(ba)^k$ are both equal to the set of all words of length k over $\{a, b\}$ provides a lower bound on k -reconstructibility. Here we can use a very similar example: the sets of k -deletions of $(12)^k$ and $(21)^k$ are both equal to the set of all compositions of $2k$ in which no entry is greater than 2. This implies that Theorem 1 is best possible.

The proof. Our reconstruction algorithm/proof of Theorem 1 employs several composition statistics. One is the *exceedance number*, defined by $\text{ex}(w) = \|w\| - |w| = \sum (w_i - 1)$ where the sum is over all entries $w(i)$. Another important composition statistic is the number of 1's in w , which can be approximated using its set of k -deletions:

Lemma 2. *The composition w of $n \geq 3k + 1$ has at least k 1's if and only if either*

- (1) 1^{n-k} is a k -deletion of w , or
- (2) the longest k -deletion of w is k letters longer than the shortest k -deletion of w .

Moreover, w has precisely k 1's if and only if one of the above conditions holds and w has a k -deletion without 1's.

Proof. It is easy to see that if either (1) or (2) occurs then w has at least k 1's. Suppose then that w has at least k 1's. If $\text{ex}(w) \leq k$ then 1^{n-k} is a k -deletion of w , satisfying (1). On the other hand, if $\text{ex}(w) > k$ then some k -deletion of w has length $|w|$, while the fact that w contains at least k 1's guarantees that some k -deletion of w has length $|w| - k$, satisfying (2). The second claim in the lemma is then readily verified. $\square$

[†]This partial order was first considered by Bergeron, Bousquet-Mélou, and Dulucq [1], and has since been studied by Snellman [12, 13], Sagan and Vatter [10], and Björner and Sagan [2].

Given a set of k -deletions of a composition, the first step in our algorithm is to apply Lemma 2 to decide if the composition has fewer than k , precisely k , or more than k 1's. The three cases are handled separately. The first two are relatively straightforward, while the last is more delicate.

Lemma 3. *If w is a composition of $n \geq 3k + 1$ with fewer than k 1's, then w can be reconstructed from its set of k -deletions.*

Proof. Given the set of k -deletions of a composition w satisfying these hypotheses, our algorithm can apply the result of Lemma 2 to determine that w has fewer than k 1's. It then follows that

$$\text{ex}(w) \geq \frac{\|w\| - (\# \text{ of 1's in } w)}{2} \geq \frac{2k + 2}{2} = k + 1.$$

From this we see that w has the same length, say m , as its longest k -deletions, and then $\text{ex}(w)$ can be easily determined: it is k plus the exceedance number of one of the longest k -deletions.

Set $t = \text{ex}(w) - k$ and define the composition $a = a(1) \cdots a(m)$ by

$$a(i) = \max\{s : \underbrace{1 \cdots 1}_{i-1} s \underbrace{1 \cdots 1}_{m-i} \text{ is, or is contained in, a } k\text{-deletion of } w\}.$$

It follows that a satisfies

$$a(i) = \min\{w(i), t + 1\}. \quad (1)$$

There are now two cases in which we are done:

- If $\|a\| = n$ then w must be equal to a . By (1), this will occur if w contains no entries greater than $t + 1$.
- If at most one entry of a satisfies $a(i) = t + 1$ — which by (1) will occur if w contains at most one entry $w(i) \geq t + 1$ — then (1) forces $w(j) = a(j)$ for all $j \neq i$ and then $w(i)$ can be calculated from the fact that $\|w\| = n$.

Suppose, for the sake of contradiction, that neither of these conditions hold. Thus w must contain an entry $w(i) > t + 1$ and another entry $w(j) \geq t + 1$. We then have

$$k + t = \text{ex}(w) \geq t + (t + 1) + (\# \text{ of } \geq 2 \text{ entries in } w, \text{ not including } w(i), w(j)),$$

so

$$k \geq t + 1 + (\# \text{ of } \geq 2 \text{ entries in } w, \text{ not including } w(i), w(j)), \quad (2)$$

while

$$|w| = 2 + (\# \text{ 1s in } w) + (\# \text{ of } \geq 2 \text{ entries in } w, \text{ not including } w(i), w(j)),$$

so because w contains fewer than k 1's,

$$(\# \text{ of } \geq 2 \text{ entries in } w, \text{ not including } w(i), w(j)) \geq |w| - k - 1. \quad (3)$$

Combining (2) and (3) shows that $|w| \leq 2k - t$, but then $\text{ex}(w) \geq (3k + 1) - (2k - t) = k + t + 1$, contradicting the definition of t and completing the proof. $\square$

Example 4. Suppose the reconstruction algorithm is given the set of 3-deletions

$$\{52, 322, 412, 421, 511, 2122, 3112, 3121, 4111\}$$

of an unknown composition w of $n = 10$. The algorithm first checks the hypotheses of Lemma 2. The first condition does not hold because the set of 3-deletions does not contain $1^{10-3} = 111111$, while the second condition fails because the longest 3-deletion is only 2 letters longer than the shortest. Therefore w has fewer than $k = 3$ 1's. Now the algorithm follows the proof of Lemma 3. First we compute $\text{ex}(w)$ from one of the longest 3-deletions:

$$\text{ex}(w) = \text{ex}(3121) + 3 = 6,$$

so $t = 3$. Then we compute a :

$$\begin{aligned} a(1) &= 4 \text{ because } 4111 \text{ is contained in a 3-deletion but } 5111 \text{ is not,} \\ a(2) &= 1 \text{ because } 1111 \text{ is contained in a 3-deletion but } 1211 \text{ is not,} \\ a(3) &= 2 \text{ because } 1121 \text{ is contained in a 3-deletion but } 1131 \text{ is not,} \\ a(4) &= 2 \text{ because } 1112 \text{ is contained in a 3-deletion but } 1113 \text{ is not.} \end{aligned}$$

Thus $w \geq 4122$. Since $\|4122\| = 9 < 10 = \|w\|$, we are not done reconstructing w and need to account for one more exceedance. However, since $a(1)$ is the only entry of a equal to $t + 1 = 4$, $w(1)$ is the only entry of w that can be greater than the corresponding entry of a , so we get $w = 5122$.

Lemma 5. *If w is a composition of $n \geq 3k + 1$ with precisely k 1's, then w can be reconstructed from its set of k -deletions.*

Proof. Given the set of k -deletions of a composition w satisfying these hypotheses, our algorithm can apply the result of Lemma 2 to determine that it has exactly k 1's. With this established, the length of w can be computed as k plus the length of the shortest k -deletion of w .

There is a k -deletion of w without 1's, and this composition gives the ≥ 2 entries of w in their correct order. Thus it suffices to determine where they lie in w . To this end define the composition a_i by

$$a_i = \underbrace{1 \cdots 1}_{i-1} 2 \underbrace{1 \cdots 1}_{m-i}.$$

As a_i is contained in a k -deletion of w if and only if $w(i) \geq 2$, the ≥ 2 entries of w can be discerned, completing the proof. $\square$

Example 6. Suppose the reconstruction algorithm is given the set of 3-deletions

$$\{322, 2212, 2221, 3112, 3121, 3211, 12121, 12211, 21121, \\ 21211, 22111, 31111, 111211, 121111, 211111\}.$$

of an unknown composition w of $n = 10$. Since the longest 3-deletions in this set are 3 letters longer than the shortest 3-deletion, w has at least $k = 3$ 1's by Lemma 2. As the set

also contains a 3-deletion without 1's, the same lemma shows that w has precisely 3 1's, and thus the algorithm follows the proof of Lemma 5. The 3-deletion without 1's — 322 — gives the ≥ 2 entries of w in their correct order. Now we form the a_i 's to see where these ≥ 2 entries lie:

$$\begin{aligned} a_1 &= 211111 \text{ is contained in a 3-deletion so } w(1) \geq 2, \\ a_2 &= 121111 \text{ is contained in a 3-deletion so } w(2) \geq 2, \\ a_3 &= 112111 \text{ is not contained in a 3-deletion so } w(3) = 1, \\ a_4 &= 111211 \text{ is contained in a 3-deletion so } w(4) \geq 2, \\ a_5 &= 111121 \text{ is not contained in a 3-deletion so } w(5) = 1, \\ a_6 &= 111112 \text{ is not contained in a 3-deletion so } w(6) = 1. \end{aligned}$$

Therefore we get $w = 321211$.

This leaves us to consider the case of compositions with many 1's. In this case we also need the *second exceedance number*, defined by $\text{ex}_2(w) = \sum (w(i) - 2)$ where the sum is over all entries $w(i) \geq 2$.

Lemma 7. *If w is a composition of $n \geq 3k + 1$ with more than k 1's, then w can be reconstructed from its set of k -deletions.*

Proof. Given the set of k -deletions of such a composition w , our algorithm can apply the result of Lemma 2 to conclude that it has more than k 1's. Therefore the k -deletions with the fewest 1's contain all ≥ 2 entries of w in the order in which they occur in w ; let $v = v(1) \cdots v(\ell)$ denote the composition formed by these entries, so

$$w = \underbrace{1 \cdots 1}_{z(1)} v(1) \underbrace{1 \cdots 1}_{z(2)} v(2) \cdots v(\ell - 1) \underbrace{1 \cdots 1}_{z(\ell)} v(\ell) \underbrace{1 \cdots 1}_{z(\ell+1)}$$

for some word $z \in \mathbb{N}^{\ell+1}$ (we take $\mathbb{N}$ to denote the nonnegative integers). Our goal is thus to determine z . We use similar techniques as in the proof of Lemma 3, although here we must perform two steps.

The first of these steps is to find the 0's in z . For $1 \leq i \leq \ell + 1$ let

$$a_i = \underbrace{2 \cdots 2}_{i-1} 1 \underbrace{2 \cdots 2}_{\ell+1-i}.$$

Since the 2's in a_i can only embed into ≥ 2 's in w , if a_i is contained in a k -deletion of w then its 1 must embed into an element between $v(i-1)$ and $v(i)$, implying that $z(i) \geq 1$. Conversely, if a_i is not contained in a k -deletion of w then either $\|a_i\| > n - k$ or $z(i) = 0$. Simple accounting shows that

$$n - k = ((\# \text{ of 1's in } w) + 2\ell + \text{ex}_2(w)) - k,$$

so $\|a_i\| = 2\ell + 1 \leq n - k$ because w has more than k 1's, and thus

$$z(i) = 0 \iff a_i \text{ is not contained in a } k\text{-deletion of } w. \quad (4)$$

The second step is to use these 0's to divine the nonzero entries of z . Define the composition $b_i = b_i(1) \cdots b_i(\ell)$ by

$$b_i(j) = \begin{cases} 1 & \text{if } j \leq i-1 \text{ and } z(j) = 0 \text{ or} \\ & j \geq i \text{ and } z(j+1) = 0, \text{ or} \\ 2 & \text{otherwise,} \end{cases}$$

and consider the possible embeddings of b_i in w . Suppose for the sake of example that $i \geq 4$. If $z(1) \geq 1$ then $b_i(1) = 2$ and thus can embed only into or to the right of $v(1)$. Otherwise if $z(1) = 0$ then $b_i(1) = 1$, but in this case $v(1)$ is the first entry of w so again $b_i(1)$ can embed only into or to the right of $v(1)$. Continuing this manner, if $z(2) \geq 1$ then $b_i(2) = 2$, and since $b_i(2)$ can only embed into a ≥ 2 entry in w to the right of $b_i(1)$, $b_i(2)$ can only embed into or to the right of $v(2)$. Otherwise if $z(2) = 0$ then $b_i(2) = 1$, but then $v(1)$ and $v(2)$ are adjacent in w so since $b_i(1)$ must embed into or to the right of $v(1)$ and $b_i(2)$ must embed to the right of $b_i(1)$ we see that $b_i(2)$ must embed into or to the right of $v(2)$. Continuing in this manner it is easy to see (or more formally, to prove inductively) that:

- For all $j \leq i-1$, $b_i(j)$ must embed into or to the right of $v(j)$.
- For all $j \geq i$, $b_i(j)$ must embed into or to the left of $v(j)$.

These two facts combine to show that $b_i(i-1)$ and $b_i(i)$ can only embed between $v(i-1)$ and $v(i)$ (inclusive). Now define the word $x \in \mathbb{N}^{\ell+1}$ by $x(i) = 0$ if $z(i) = 0$ and otherwise

$$x(i) = \max\{s : b_i(1) \cdots b_i(i-1) \underbrace{1 \cdots 1}_s b_i(i) \cdots b_i(\ell) \text{ is contained in a } k\text{-deletion of } w\}.$$

The analogue to (1) now follows by the conditions on embeddings of b_i established above:

$$x(i) = \min\{z(i), n - k - \|b_i\|\}. \quad (5)$$

Suppose $z(i) \geq 1$. In this case $\|b_i\| = 2\ell - h$, where h denotes the number of 0 entries of z ("holes"). Letting $k + t$ denote the number of 1's in w , we have

$$n = k + t + 2\ell + \text{ex}_2(w).$$

allowing us to rewrite (5) as

$$x(i) = \min\{z(i), h + t + \text{ex}_2(w)\}. \quad (6)$$

If $\|v\| + \|x\| = n$ then we must have $z = x$ and thus have successfully reconstructed w . By (6), this will happen if z has no entries greater than $h + t + \text{ex}_2(w)$. Suppose, for the sake of contradiction, that this does not occur, i.e., that z contains an entry greater than $h + t + \text{ex}_2(w)$. Then each of the other $(\ell + 1 - h) - 1$ nonzero entries of z correspond to at least one 1 in w , and thus we have

$$k + t = \# \text{ of 1's in } w \geq (h + t + \text{ex}_2(w) + 1) + (\ell - h) = t + \ell + \text{ex}_2(w) + 1.$$

However, this implies that

$$2k \geq t + 2\ell + \text{ex}_2(w),$$

so

$$3k \geq (k + t) + 2\ell + \text{ex}_2(w) = n,$$

and this contradiction completes the proof of both the lemma and Theorem 1. $\square$

Example 8. Suppose the reconstruction algorithm is given the set of 3-deletions

$$\{1222, 2212, 11122, 11212, 11221, 12112, 12211, 111112, 111121, 111211, 112111, 1111111\}.$$

of an unknown composition w of $n = 10$. This set contains $1^{10-3} = 1111111$ and every 3-deletion in the set contains a 1, so Lemma 2 shows that w has more than $k = 3$ 1's. Thus we follow the proof of Lemma 7. Each of the compositions with the fewest 1's, e.g., 2122, give the ≥ 2 entries of w in their correct order, $v = 222$, so

$$w = \underbrace{1 \cdots 1}_z 2 \underbrace{1 \cdots 1}_z 2 \underbrace{1 \cdots 1}_z 2 \underbrace{1 \cdots 1}_z.$$

We then find the 0 entries of z :

- $z(1) \neq 0$ because $a_1 = 1222$ is contained in a 3-deletion of w ,
- $z(2) = 0$ because $a_2 = 2122$ is not contained in a 3-deletion of w ,
- $z(3) \neq 0$ because $a_3 = 2212$ is contained in a 3-deletion of w ,
- $z(4) = 0$ because $a_4 = 2221$ is not contained in a 3-deletion of w .

Now we build the word $x \in \mathbb{N}^4$. We have that $x(2) = x(4) = 0$ because the corresponding entries of z are 0. To compute the other entries of x we construct $b_1 = 121$ and $b_3 = 211$ and then have

- $x(1) = 3$ because 111 121 is contained in a 3-deletion of w but 1111 121 is not,
- $x(3) = 1$ because 21 1 1 is contained in a 3-deletion of w but 21 11 1 is not.

Since $\|v\| + \|x\| = \|222\| + \|3010\| = 10$, we must have $z = x$ and thus $w = 1112212$.

The connection to permutations. The subject of permutation patterns (see Bóna's text [3] for a survey) is concerned with the following partial order on permutation: for permutations σ of length k and π of length n , let $\sigma \leq \pi$ if there are indices $i_1 < i_2 < \cdots < i_k$ such that the subsequence $\pi(i_1)\pi(i_2)\cdots\pi(i_k)$ has the same pairwise comparisons as $\sigma(1)\sigma(2)\cdots\sigma(k)$, and in such a case σ is said to be an $(n - k)$ -deletion of π . For example, $13254 \leq 213654798$ because of the subsequence $26598 (= \pi(1)\pi(4)\pi(5)\pi(8)\pi(9))$.

Given two permutations σ and π of lengths m and n respectively, their *direct sum*, $\sigma \oplus \pi$, is the permutation of length $m + n$ whose first m entries form σ and whose last n entries are the copy of π obtained by adding m to each entry. For example, $213654 \oplus 132 = 213654798$.

A permutation is said to be *layered* if it can be written as the direct sum of decreasing permutations. Thus 213654798 is layered because it can be written as $21 \oplus 1 \oplus 321 \oplus 1 \oplus 21$. There is a natural order-preserving bijection between layered permutations and compositions; for example, $213654798 = 21 \oplus 1 \oplus 321 \oplus 1 \oplus 21$ maps to the composition 21312 while $13254 = 1 \oplus 21 \oplus 21$ maps to 122, and $122 \leq 21312$ under the partial order on compositions.

Smith [11] was the first to study multiset reconstruction for permutations. Her work was followed by Raykova [9] who proved that for all k , all sufficiently long permutations are reconstructible from their multisets of k -deletions. This leaves open the question of whether all sufficiently long permutations are reconstructible from their *sets* of k -deletions. Our work therefore answers Raykova's question about whether all sufficiently long layered permutations can be reconstructed from their sets of k -deletions.

Acknowledgement. I thank Robert Brignall for his helpful comments.

REFERENCES

- [1] BERGERON, F., BOUSQUET-MÉLOU, M., AND DULUCQ, S. Standard paths in the composition poset. *Ann. Sci. Math. Québec* 19, 2 (1995), 139–151.
- [2] BJÖRNER, A., AND SAGAN, B. Rationality of the Möbius function of a composition poset. *Theoret. Comput. Sci.* 359, 1-3 (2006), 282–298.
- [3] BÓNA, M. *Combinatorics of permutations*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2004. With a foreword by Richard Stanley.
- [4] CAMERON, P. J. Stories from the age of reconstruction. *Congr. Numer.* 113 (1996), 31–41.
- [5] HARARY, F. On the reconstruction of a graph from a collection of subgraphs. In *Theory of Graphs and its Applications (Proc. Sympos. Smolenice, 1963)*. Publ. House Czechoslovak Acad. Sci., Prague, 1964, pp. 47–52.
- [6] LOTHAIRE, M. *Combinatorics on words*, vol. 17 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Co., Reading, Mass., 1983.
- [7] MANVEL, B. Some basic observations on Kelly's conjecture for graphs. *Discrete Math.* 8 (1974), 181–185.
- [8] PRETZEL, O., AND SIEMONS, J. On the reconstruction of partitions and applications. *Electron. J. Combin.* 11 (2) (2005), Note 5, 6 pp. (electronic).
- [9] RAYKOVA, M. Permutation reconstruction from minors. *Electron. J. Combin.* 13 (2006), Research paper 66, 14 pp. (electronic).

- [10] SAGAN, B. E., AND VATTER, V. The Möbius function of a composition poset. *J. Algebraic Combin.* 24, 2 (2006), 117–136.
- [11] SMITH, R. Permutation reconstruction. *Electron. J. Combin.* 13 (2006), Note 11, 8 pp. (electronic).
- [12] SNELLMAN, J. Saturated chains in composition posets. arXiv:math.CO/0505262.
- [13] SNELLMAN, J. Standard paths in another composition poset. *Electron. J. Combin.* 11, 1 (2004), Research Paper 76, 8 pp. (electronic).